Survey on Dermatophytosis in Iraqi camels

M. H. Hussain

Col. of Vet. Med./ Al-Qadissiya Univ.

Abstract

This survey aimed to answer a question has applied in the clinics “what are the fungal skin affections in Iraqi camels?”. The survey, in three provinces; Al-Qadissiya, Al- Najaf & Al-Muthanna; has identified three Dermatophytes “*Trichophyton verrucosum* in 58%, *Microsporum canis* in 42% & *Epidermophyton fluccosum* in 22%” & four non-dermatophytes “*Penicillium brevicompactum* in 31.6%, *Ulocladium chartarum* in 22.8%, *Aspergillus fumigates* in 22.8% & *Scopulariopsis bainier* in 22.8%” in 112 camels. Dermatophytosis has diagnosed in 65.2% in camels less than 5 years old, 34.8% in camels 5-10 years old & no cases were diagnosed in camels more than 10 years old.

Introduction

Dermatophytosis is an infection of keratinized tissue (skin, hairs, and claws) by one of the three genera of fungi collectively called dermatophytes; *Trichophyton*, *Microsporum*, and *Epidermophyton*. These pathogenic fungi are found worldwide, and all domestic animals are susceptible. Various fungal species can produce infection of the epidermis, of which the species causing dermatophytosis (ringworm) which are the commonest in camels. Dermatophytosis is a

common skin disease in old world camels (OWC) under 3 years of age with a peak incidence age of between 3 to 12 months. In new world camels (NWC) it is, however, a very rare disease (5) and only *T. verrucosum* and *T. mentagrophytes* have been isolated from NWC so far. In most circumstances, dermatophytes grow only in dead, keratinized tissue; advancing infection halts upon reaching live cells or inflamed tissue (11).

Materials and Methods

- 1- Potassium hydroxide sol. 10%
- 2- Sodium hydroxide sol. 10%
- 3- Lacto phenol cotton blue stain
- 4- Yeast extract
- 5- Thiamine
- 6- Chloramphenicol
- 7- Amikacin
- 8- Sabouraud dextrose agar

Direct microscopic examination of hairs or skin scrapings might reveal characteristic hyphae and/or arthrospores. However, fungal culture is the most effective and specific means of diagnosis, although growth usually requires 10 to 14 days of incubation (8). *Microsporum* spp. and *Trichophyton* spp. as well as other fungi should be cultured on Sabouraud dextrose agar and on Mycoline agar slide (bioMandieux) and incubated for 10 to 14 days at 27°C. Definite diagnosis and species identification requires removal of hyphae and macroconidia from the surface of the

colony with acetate tape and microscopic examination with Lactophenol Cotton Blue (LPCB) stain. Culture on Mycoline agar slide is especially helpful when saprophytic contamination is expected (11).

Modified Sabouraud Dextrose Agar (SDA) was prepared as below:

65 gm	Sabouraud dextrose agar
5 gm	Yeast extract
0.05 gm	Chloramphenicol
0.5 gm	Cycloheximide
0.01gm	Thiamin
1000 ml	Distilled water

Lactophenole cotton blue stain was prepared in the laboratory using:-

20 ml	Lactic acid
20 gm	Phenol crystals
40 ml	Glycerin
0.05 gm	Anelin blue
20 ml	Dist. water

Phenol crystals to be dissolved in lactic acid and water by slow heating, Anelin blue was added and mixed well (1).

Three governorates were involved in this research Al-Qadisiya, Al-Najaf and Al-Muthanna, samples collected from the slaughter houses, sale yards and the herds found in several locations randomly. The research began on December 2008 and finished on June 2009. 2412 Iraqi dromedaries were examined regardless to location, age, sex and breed, from which 482 camels were suffering skin affections. Camels were inspected to find infections with fungal lesions. Skin scrapings were collected after washing the lesion with 70% ethanol to be transported to the laboratory by using screw cap containers. (10).

Steps of identification include:-

1- Clinical signs : - lesions were prescribed according to the site, shape, color,

hair coat and any other signs. The general description of camels including; temperature, respiratory rate, pulse rate, mucous membrane, appetite and ruminating and other features.

2- Scrapings were taken to be cultured on the Modified Sabouraud Dextrose Agar SDA that was prepared in the laboratory. For purification, multiple cultures were done to obtain clear and pure colonies without contaminations. Media were incubated at 28°C for up to 2 weeks according to the growth appearance. Each type of dermatophytes were examined microscopically by making slide smears from the pure colonies and stained with Lactophenole Cotton Blue Stain LCB. Different characteristic features confirmed the diagnosis which depends on the shape of the colony, the microscopic appearance and microscopic examinations of the hides.

Results

The survey has found 112 camels infected with dermatophytosis in different sexes but in less than 10 years old and most commonly in camels less than 5 years old with 65.2% and 34.8% in camels in 5-10 years old. It seems that camels less than 5 years old are more susceptible to fungal infection with dermatophytes. Both sexes are likely to have the same percentage of infection, as in table (2). Three dermatophytes and four non-dermatophytes have been diagnosed in the laboratory after

sampling skin scrapings from the suspected camels as illustrated in table (1), the diagnosis depends on the duration of growth, features of the colonies and the microscopic appearance after preparing the slides stained with the lactophenole cotton blue which is the most effective and specific means for fungal diagnosis (6) and (7). As showed in table (1), seven species of fungi were described according to the duration of growth in the colonies obtained after culture.

Table (1): Dermatophytes were diagnosed in the study.

Dermatomycosis		duration of growth (day)
Dermatophytes	<i>Trichophyton verrucosum</i>	10-14
	<i>Microsporum canis</i>	8-12
	<i>Epidermophyton fluccosum</i>	4-7
Non-dermatophytes	<i>Penicillium brevicompactum</i>	2-5
	<i>Ulocladium chartarum</i>	3-5
	<i>Aspergillus fumigates</i>	2-4
	<i>Scopulariopsis bainier</i>	3-6

Trichophyton verrucosum was the most common case 58% accompanied with *Epidermatophyton fluccosum* in 9 cases, the lesions were multifocal on the neck, legs, head, shoulders, chest, and perineum respectively in which, as in fig (1), gray-white crusts raised slightly on the skin in circles within 1-3 cm in diameter with moist surface beneath in the early stage losing hair, when the lesion over growth the surface become hairless, gray, dry and the crusts detached and the circles conjoin to present the whole site on the infected area. It seems to be due to the slow spread of lesions in the herd and their self-limiting in about 25-40 days with normal skin finally. *Microsporum*

canis have the same clinical features, but the lesions spreaded mostly in the perineum and they looked irregular shapes and the spots unite soon after with rapid hair loose, *E. fluccosum* had diagnosed in 16 camels accompanied with *M. canis* on the same individuals. No systemic reactions to be observed on the camels having ring worm, (36.3 ± 0.03) C° in temperature, (15 ± 0.5) respiratory rate, (45 ± 1.5) pulse rate, pink mucous membrane, hair loss, no lameness when the leg infected, normal appetite and ruminating and itching seems to be rare. Accidental respiratory disorders or diarrhea have seen in several cases.

Table (2): Numbers of infected camels with cutaneous fungi according to the location, age and sex

Locations		Infected camels	~5 years		5-10 years		10~ years	
			male	female	male	female	male	female
Al-Qadissyya	Slaughter house	19	1	10	5	3	0	0
	Al-Hamza Al-Shargy	6	3	1	0	2	0	0
	Al-Daghara	0	0	0	0	0	0	0
	Aal-Bdeer	11	2	5	2	2	0	0
	Al-Shaf'ya	0	0	0	0	0	0	0
	Al-Shenafeya	16	3	8	1	4	0	0
Al-Najaf	Slaughter house	19	11	2	3	3	0	0
	Sale yard	22	9	4	6	3	0	0
Al-Muthanna	Slaughter house	2	2	0	0	0	0	0
	Sale yard	0	0	0	0	0	0	0
	Herds	17	3	9	2	3	0	0
Total		112	34	39	19	20	0	0
Percentage			30.4	34.8	17	17.8	0	0
Total			73		39		0	
Percentage			65.2		34.8		0

Fig (1): *Trichophyton verrucosum* (ring worm) in 3.5 years camel, circular lesions on the neck.

Colony of the *T. verrucosum* on the modified SDA slowly appears heaped, white powdery surface at the first few days; within incubation the center becomes brown illustrated in fig (2). Microscopically, slide smear stained with lactophenol cotton blue

shows, as in fig (3), thin irregular mycelium with chlamydospores that form heavy chains. Macroconidia are extremely rare, three to five celled with thin and small walls vary considerably in size and shape.

Fig (2): Colony of the *T. verrucosum* on the modified SDA.

Fig (3): Microscopic appearance of *T. verrucosum*.X100

Discussion

Camels' dermatophytosis had been reported in Iraq by Al-Ani, *et al.*, 1995 through a study found out infections with *Trichophyton schoenleinii*. 13 non-dermatophytes reported in Al-Basrah by Abdul-Aziz (2000), only *A.fumigatus* was corporate and the other types were different. Differences may belong to the variance between Al-Qadisiya and Al-Basrah in several factors; type of the soil, climatic features, temperature and humidity, regarding to the fact that several non-dermatophytes are soil-borne fungi while others may be food-borne fungi. Consequently, there will be some differences in the types of fungi that suspended to

animals' skin. Absence of infections in camels older than 10 years may reveals the formation of immunity against this cutaneous pathogen, but also it has reported that ring worm is more common in camels less than 5 years old. Other types of dermatophytes had reported in the world, but it's obvious, that *T. verrucosum* is the most vulgar dermatophyte since it has reported by Curasson in 1947. Diagnosis was made by the help of Introduction of food-borne fungi (9) and the Introducing mycology by examples, texts with pictures for diagnosis of fungi, a special help from Prof. Dr. M. K. Faraj, Dept of Microbiology/ College of Veterinary Medicine, Baghdad University.

References

- Pathogenic Fungi. Your Book Medical Publishers. Chicage. 268.
- 7- Midgley, G.; Clayton, Y. and Hay, R. J. (1997). Diagnosis in color medical mycology Mosby Wolfe, an imprint of Mosby international. Spain. 155.
- 8- Radostits, O.M.; Gay, C.C.; Hinchcliff, K.W.; & Constable, P.D. (2007). Veterinary Medicine, A textbook of the diseases of cattle, horses, sheep, pigs & goats. 10th ed. Elsevier Saunders, Spain.
- 9- Rebert, A.S.; Ellen, S.H.; & Connie, A.N.(1984). Itroduction of food-borne fungi. 2nd, Central bureau voor schimmel cultures. Institute of the royal Netherl&s, Academy of art & science.
- 10- Songer, J.G.; & Post, K.W.(2005). Veterinary Microbiology, Bacterial & Fungal agents of animal diseases, chapter four; the genus Staphylococcus. Elsevier Saunders, china.
- 11- Wernery, U.; & Kaaden, O.R. (2002). Infectious diseases in camelids, 2nd revised & enlarged edition. Blackwell Wissenschafts-Verlag
- 1- حسن, وهاب امين و الخطيب, غازي موسى (1990). مبادئ الاحياء المجهرية, ملحق 3: بعض الاصباغ المستعملة في الفطريات ص 533-536, هيئة المعاهد الفنية/ وزارة التعليم العالي والبحث العلمي, مطابع دار الحكمة للطباعة والنشر.
- 2- عبد العزيز, جمال مهدي (2000). دراسة حول الإصابات الفطرية في الأبقار والفطريات العالقة بشعر الأبقار والجاموس والخيول والإبل في محافظة البصرة وتأثير خمسة أنواع من المضادات الفطرية على بعض العزلات. رسالة ماجستير / كلية العلوم / جامعة البصرة.
- 3- Al-Ani, F. K.; Al-Bassam, L.S. & Al-Salahi, K.A. (1995). Epidemiological study of dermatomycosis due to *Trichophyton schoenleinii* in camels in Iraq. *Bull. Anim. Hlth. Prod. Afi.* 43: 87-92.
- 4- Curasson, G. (1947). Le chameau et ses maladies. Vigot Fr&es, Editeurs: pp. 86-88.
- 5- Fowler, M.E. (1998).Medicine & surgery of South American Camelids. Iowa State University Press, Ames. USA.
- 6- Frey, D.; Oldfield, R. J. and Bridger, R. C. (1979). A Color Atlas of
- 12- GmbH Kurfiirstendamm 57,10707 Berlin

مسح ميداني عن الفطريات الجلدية في الإبل العراقية

مثنى هادي حسين *

كلية الطب البيطري/ جامعة القادسية

الخلاصة

إجابة على السؤال الذي طالما يطرح في العيادات والمستشفيات البيطرية "ما هي الإصابات الجلدية الفطرية في الإبل العراقية؟" جاءت هذه الدراسة أو المسح الميداني الذي شمل ثلاث محافظات وهي القادسية، النجف والمثنى لتكشف عن وجود ثلاث أنواع من الفطريات الجلدية الممرضة وهي:

Trichophyton verrucosum in 58%.

Microsporum canis in 42%.

Epidermophyton fluccosum in 22%.

بالإضافة إلى أربع أنواع من الفطريات الجلدية غير الممرضة وهي:

Penicillium brevicompactum in 31.6%.

Ulocladium chartarum in 22.8%.

Aspergillus fumigates in 22.8%.

Scopulariopsis bainier in 22.8%.

تم تشخيص الإصابات الفطرية الجلدية في الإبل دون سن الخامسة بنسبة 65.2%, وبنسبة 34.8% في الإبل بين سن الخامسة والعاشرة في حين لم تسجل أي إصابة عند الإبل في الأعمار الأكبر من عشرة سنوات.

* البحث مستل من أطروحة الماجستير للباحث.